



ORIGINAL ARTICLE

Validity and Reliability Study of the Korean Tinetti Mobility Test for Parkinson's Disease

Jinse Park,¹ Seong-Beom Koh,² Hee Jin Kim,³ Eungseok Oh,⁴ Joong-Seok Kim,⁵
Ji Young Yun,⁶ Do-Young Kwon,⁷ Younsoo Kim,⁸ Ji Seon Kim,⁹
Kyum-Yil Kwon,¹⁰ Jeong-Ho Park,¹¹ Jinyoung Youn,^{12,13} Wooyoung Jang¹⁴

¹Department of Neurology, Haeundae Paik Hospital, Inje University, Busan, Korea

²Department of Neurology & Parkinson's Disease Centre, Guro Hospital, Korea University, Seoul, Korea

³Department of Neurology, Konkuk University School of Medicine, Seoul, Korea

⁴Department of Neurology, Chungnam National University Hospital, Daejeon, Korea

⁵Department of Neurology, Seoul St. Mary's Hospital, The Catholic University of Korea, Seoul, Korea

⁶Ewha Womans University School of Medicine, Ewha Womans University Mokdong Hospital, Seoul, Korea

⁷Department of Neurology, Korea University College of Medicine, Ansan Hospital, Ansan, Korea

⁸Department of Neurology, Samsung Changwon Hospital, Sungkyunkwan University School of Medicine, Changwon, Korea

⁹Department of Neurology, Chungbuk National University College of Medicine, Cheongju, Korea

¹⁰Department of Neurology, College of Medicine, Soonchunhyang University Seoul Hospital, Seoul, Korea

¹¹Department of Neurology, College of Medicine, Soonchunhyang University Bucheon Hospital, Bucheon, Korea

¹²Department of Neurology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

¹³Neuroscience Center, Samsung Medical Center, Seoul, Korea

¹⁴Gangneung Asan Hospital, University of Ulsan College of Medicine, Gangneung, Korea

ABSTRACT

Objective Postural instability and gait disturbance are the cardinal symptoms associated with falling among patients with Parkinson's disease (PD). The Tinetti mobility test (TMT) is a well-established measurement tool used to predict falls among elderly people. However, the TMT has not been established or widely used among PD patients in Korea. The purpose of this study was to evaluate the reliability and validity of the Korean version of the TMT for PD patients.

Methods Twenty-four patients diagnosed with PD were enrolled in this study. For the interrater reliability test, thirteen clinicians scored the TMT after watching a video clip. We also used the test-retest method to determine intrarater reliability. For concurrent validation, the unified Parkinson's disease rating scale, Hoehn and Yahr staging, Berg Balance Scale, Timed-Up and Go test, 10-m walk test, and gait analysis by three-dimensional motion capture were also used. We analyzed receiver operating characteristic curve to predict falling.

Results The interrater reliability and intrarater reliability of the Korean Tinetti balance scale were 0.97 and 0.98, respectively. The interrater reliability and intra-rater reliability of the Korean Tinetti gait scale were 0.94 and 0.96, respectively. The Korean TMT scores were significantly correlated with the other clinical scales and three-dimensional motion capture. The cutoff values for predicting falling were 14 points (balance subscale) and 10 points (gait subscale).

Conclusion We found that the Korean version of the TMT showed excellent validity and reliability for gait and balance and had high sensitivity and specificity for predicting falls among patients with PD.

Key Words Parkinson's disease; Tinetti mobility test; gait; balance.

Received: August 21, 2017 Revised: October 16, 2017 Accepted: October 30, 2017

Corresponding author: Seong-Beom Koh, MD, Department of Neurology & Parkinson's Disease Centre, Guro Hospital, Korea University, 148 Guro-dong-ro, Guro-gu, Seoul 08308, Korea / Tel: +82-2-2626-1114 / Fax: +82-2-797-0298 / E-mail: parkinson@korea.ac.kr

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gait disturbance and postural instability are cardinal features of Parkinson's disease (PD) and detract from quality of life. Objective measurement of gait and balance to observe the progression of disease and reduce the fall risk of patients has been the subject of much attention.¹ Most clinicians assess gait and balance by relying on subjective observations during the neurological examination. Several clinical scales for gait and balance have been developed. The most commonly used scales for gait and balance status in PD are Hoehn and Yahr (H&Y) staging and the Unified Parkinson's Disease Rating Scale (UPDRS) part III.^{2,3} H&Y staging is a simple and well-known method; however, the H&Y scale is not linear and cannot even be ordered by rank.⁴ The UPDRS part III contains items of gait and postural instability, but the scoring is also broad and simple. The Berg balance test, which is also widely used in estimating postural imbalance, has been validated for PD.⁵ However, the Berg balance test only can estimate balance and cannot evaluate gait itself.

As a way overcome these limitations of clinical scales, the Tinetti Mobility Test (TMT), also called the Performance-Oriented Mobility Assessment, was introduced.⁶ The TMT has been widely used and has been recommended as a way to assess mobility, balance, gait, and fall risk in the elderly.⁷ Previous reports have demonstrated that the TMT is one of the most accurate tools for assessing patients with PD and has acceptable validity and reliability.⁸ The TMT was translated into Korean in a previous study and was shown to be a reliable and valid tool for chronic stroke patients.⁹ However, there has been no report of the validation of the Korean version of the TMT in PD patients.

The purpose of this study was to investigate the reliability and validity of the Korean version of the TMT in patients with PD.

MATERIALS & METHODS

This is a cross-sectional observational study. The primary goal was to evaluate the reliability and validity of the Korean version of the TMT in patients with PD. The secondary goal was to identify the most accurate TMT cutoff value for predicting falls in patients with PD. This study was approved by Interstitial Review Board in Haeundae Paik Hospital (IRB No. 2016-04-024).

Subjects

Twenty-four patients diagnosed with PD according to the Queen Square Brain Bank criteria¹⁰ at three different hospitals were enrolled in this study. The inclusion criteria were as follows: 1) PD patients who complained of gait disturbance and postural instability, 2) patients with PD aged 50 to 80 years, 3) H&Y stage 2 to 4. All participants were given informed consent.

The exclusion criteria were as follows: 1) patients who were diagnosed with dementia, 2) patients who had an evident risk of falling while performing the test, 3) patients who had other diseases affecting gait such as neuromuscular or cardiopulmonary dysfunction, and 4) patients who were diagnosed with atypical parkinsonism.

Basic demographic factors including age, sex, disease duration, and falling history were obtained. We used the UPDRS motor scale (part III), the balance subscale (item 30), and the H&Y stage to evaluate disease severity. We obtained each patient's falling history from an interview at baseline screening. A fall was defined as an unexpected falling event that ended with the subject coming to rest on the ground and was not due to an extrinsic event. A "faller" was defined as a person who had fallen more than once within six months.¹¹ We categorized the patients into faller and non-faller groups on the basis of their falling history.

Korean version of the TMT

Although a Korean TMT has been published for stroke patients, this published Korean TMT has several limitations: 1) the paper introducing the Korean TMT is written in Korean, which makes it difficult to find through online searches in PubMed or Embase; 2) the published Korean TMT includes unfamiliar term such as stance phase that are difficult to understand by the public. 3) only two examiners were used to estimate interrater reliability. Therefore, we constructed a new version of the Korean TMT according to the following method.

The TMT as tested in this study is composed of two distinct components, including a balance subscale (9 items, 16 points) and a gait subscale (8 items, 12 points). Therefore, there are 17 items, each of which is rated on a scale of 0 to 1 or 2, and the maximum possible score is 28 points. The total time taken to perform the TMT is approximately 10 minutes

per participant. The translation process of the TMT is accurate and reflects Korean culture.

For the translation of the TMT into its Korean version, one movement specialist and one non-medical-expert translator participated in the translation. Subsequently, another non-medical-expert translator retranslated the Korean TMT into the English version and compared it with the original version of the TMT. Each non-medical-expert translator was a person not associated with healthcare who had lived in an English-speaking country for at least 10 years. The three people who participated in the translation collaborated to adapt it to Korean culture (Supplementary Table 1 in the online-only Data Supplement).

Checking the reliability and validity of the TMT

Three movement specialists from different hospitals discussed how to perform the TMT. All TMTs were recorded with a video camcorder with antero-posterior and mediolateral views.

For the intrarater reliability test, we used the test-retest method. We performed the TMT twice with a 30-minute interval between tests. For interrater reliability, thirteen clinicians who worked at different hospitals gathered together and scored the TMT while watching the video clips.

For concurrent validation, clinical tests for gait and balance and three-dimensional motion captures were performed. Clinical tests for gait included the Timed-up and go (TUG) test, 10-m walk test, and UPDRS part III. Clinical scales for balance and the Berg balance test were performed. For objective quantification of gait, we checked spatiotemporal parameters from the three-dimensional motion analysis.

Gait analysis by three-dimensional motion capture

To evaluate gait objectively, we used three-dimensional motion capture VICON (Oxford, UK) at a sample rate of 100 Hz for quantification of spatiotemporal parameters. The participants were asked to walk along a 6-m track under 12 infrared cameras for motion capture. Stride length, walking speed, cadence, step time, stride time, step length, single support, double support and proportion of the gait cycle spent in the stance phase were measured.

Statistics

All results were calculated using SPSS version 21 (IBM Corp., Armonk, NY, USA) and R version 3.1.2. For intra- and interrater reliability, an Interclass Correlation Coefficient (ICC) derived from a two-way random-effects model was used. For internal consistency, we calculated Cronbach's alpha in each subscale ("irr", "psy", "boot" packages).

The concurrent validity of the TMT was obtained by calculating the correlation with the Berg balance test, the UPDRS balance and gait subscores, the H&Y stage, the TUG, and the 10-m walk test. Gait parameters from three-dimensional motion analyses were also compared with TMT results. Spearman correlation analysis was used to evaluate the correlation between TMT and other parameters. The level of significance was set at $p < 0.05$.

Receiver Operating Characteristic (ROC) analysis was performed to assess the sensitivity and specificity of the TMT score in predicting falls. An ROC curve analysis and an estimate of the area under the ROC curve (AUC) were used to assess the cutoff points for TMT in PD.

RESULTS

The demographics of the participants are shown in Table 1. Their mean age was 72.33 ± 6.95 years, and fourteen of the participants were men. The mean H&Y stage was 2.39 ± 7.37 , and the mean UPDRS part III score was 15 ± 3.20 . Among the 24 participants, 8 patients had a history of falling.

Reliability of the TMT

The results of the reliability tests for the TMT are

Table 1. Demographic factors of participants

Demographic data	
M/F (n = 24)	14/10
Age	72.33 ± 6.95
UPDRS part III (total)	15 ± 3.20
Gait (item 29)	1 ± 0.20
Balance (item 30)	1 ± 0.20
H&Y stage	2.39 ± 7.37
Disease duration (months)	34.25 ± 38.23
TUG (sec)	44.25 ± 9.25
Berg balance test	22.89 ± 13.89
10-m walking (sec)	18.60 ± 13.91
Faller/non-faller (number)	8/16

UPDRS: Unified Parkinson's Disease Rating Scale, H&Y: Hoehn and Yahr, TUG: Timed-up and go.

Table 2. Reliability of the Tinetti gait and balance tests

Tinetti mobility test	Interrater reliability	Intrarater reliability	Correlation*	
			r	p
Tinetti balance scale	0.97	0.97	0.7	0.002
Tinetti gait scale	0.94	0.96	0.664	0.004

ICC derived from a two-way random-effects model is used as an estimate of raters' agreement. The interrater reliability values of the Tinetti balance and gait scales were 0.97 (0.94–0.98) and 0.94 (0.90–0.97). *mean of bivariate Pearson's product moment correlation analysis. ICC: Interclass Correlation Coefficient.

Table 3. Concurrent validity of TMT with other clinical and laboratory parameters

Clinical and laboratory scales	Tinetti balance score		Tinetti gait score	
	r	p	r	p
Berg balance test	0.73	0.00*	0.64	0.00*
TUG test	-0.64	0.00*	-0.69	0.00*
10-m walk test	-0.52	0.01*	-0.56	0.00*
UPDRS III				
Gait (item 29)	-0.45	0.03*	-0.47	0.02*
Balance (item 30)	-0.17	0.42	-0.17	0.42
H&Y staging	-0.58	0.00*	-0.38	0.06
Three-dimensional motion capture				
Stride length	0.64	0.00*	0.77	0.00*
Walking speed	0.55	0.01*	0.71	0.00*
Cadence	0.00	0.99	-0.02	0.99
Step length	0.42	0.04*	0.81	0.00*
Stance phase	-0.08	0.70	-0.74	0.00*
Stride time	0.03	0.87	-0.05	0.81
Double support	0.02	0.90	-0.56	0.00*
Single support	0.00	0.99	0.45	0.04*

*p < 0.05. TMT: Tinetti mobility test, TUG: Timed-up and go, UPDRS: Unified Parkinson's Disease Rating Scale, H&Y: Hoehn and Yahr.

shown in Table 2. The interrater reliability of the balance subscales ranged from 0.94 to 0.98 with an ICC of 0.97. The interrater reliability of the gait scale ranged from 0.90 to 0.97 with an ICC of 0.94. Test-retest reliability, used as a measure of intrarater reliability, was ICC = 0.97 for the balance scale and 0.96 for the gait scale. Cronbach's alpha coefficients for the internal consistency of the Korean version of the Tinetti balance and gait scale were 0.74 and 0.72, respectively, with a confidence interval from 0.69 to 0.77 and from 0.67 to 0.76.

Concurrent validity of Korean TMT

Concurrent validity compared with each clinical test is shown in Table 3. The Tinetti gait scale showed a negative correlation with the 10-m walking test ($r = -0.56$) and the TUG ($r = -0.64$). Regarding three-dimensional gait analysis, the Tinetti gait scale showed a positive correlation with stride length ($r = 0.77$), walking speed ($r = 0.71$), step length ($r = 0.81$) and single support ($r = 0.45$), and a negative correlation with stance phase and double support. The Tinetti balance scale showed a positive correlation with the

Table 4. ROC analysis of Tinetti balance and gait scores

Tinetti mobility test	Estimate	Lower	Upper
Balance			
Cutoff score	14		
Sensitivity	0.813	0.544	0.960
Specificity	0.750	0.349	0.968
Positive predictive value	0.867	0.538	0.973
Negative predictive value	0.667	0.355	0.953
Gait			
Cutoff score	10		
Sensitivity	0.875	0.617	0.984
Specificity	0.625	0.245	0.915
Positive predictive value	0.824	0.476	0.977
Negative predictive value	0.714	0.365	0.942

ROC: Receiver Operating Characteristic.

Berg balance scale ($r = 0.73$) and a negative correlation with the TUG ($r = -0.64$).

ROC analysis to predict falling

Table 4 shows the outcomes of the ROC analysis. The Tinetti gait scale shows high sensitivity (88%) and specificity (63%) when 10 is selected as the cutoff value. The Tinetti balance scale score of 14 was selected as the cutoff with the highest sensitivity (81%)




Figure 1. Receiver Operating Characteristic curve of Tinetti gait and balance scale for predicting falling in Parkinson's disease.

and specificity (75%) (Figure 1). The value of the AUC for predicting falls was 0.773 for the balance scale and 0.766 for the gait scale.

DISCUSSION

Our study is the first report of the reliability and validity of the TMT for evaluating patients with PD in Korea. Furthermore, we used a number of tests, including objective estimation via three-dimensional motion capture, for concurrent validation. We found that the Korean version of the TMT showed excellent reliability and validity for patients with PD. Our results are in accordance with previous reports of its high reliability and validity for patients with PD.⁸ A kappa coefficient from 0.61 to 0.80 is regarded as “substantial” reliability, and a value above 0.8 is regarded as “good” reliability.¹² In our results, the Korean version of the TMT showed excellent inter-rater and intrarater reliability. Our results showed higher inter- and intra-reliability and internal consistency than previously reported for the original TMT. Previous reports investigated its reliability for use by students and therapists, but our study targeted movement experts.¹³

The Tinetti balance scale showed a positive correlation with clinical and laboratory balance scales. Previous studies have supported a correlation between the TMT and the Berg balance test, TUG, 10-m walking test, and H&Y stage, consistent with our results.^{8,14} The Tinetti balance scale is most high-

ly correlated with the Berg balance test and TUG. The Berg balance test is the most accurate scale for estimating balance.¹¹ Interestingly, the Tinetti balance scale score did not correlate with the UPDRS balance subscore (item 30) but was significantly correlated with the Tinetti gait scale score (item 29). One possible explanation is that postural imbalance in PD is episodic and inconsistent and is therefore difficult to evaluate in one trial. However, gait performance is a constant process, and balance is needed for gaiting. As shown in Table 3, gait and postural instability are closely related to each other. The measurement of balance using the UPDRS score has limitations for evaluating balance because it was checked only by the pull test.

The Tinetti gait scale also showed a strong correlation with the Berg balance test, TUG, 10-m walk test stride length, and walking speed. Objective gait parameters including stride length and walking speed showed the strongest correlation with the Tinetti gait and balance scale. A previous study demonstrated that the Tinetti gait scale was also significantly correlated with the measurements of the GAITRite (CIR system Inc., Franklin, NJ, USA) electronic system.¹⁵ Our results demonstrated that the Korean version of the TMT was more strongly associated with gait speed and stride length than was previously reported. The 10-m walk test showed a significant but relatively weak correlation with the Korean version of the TMT.

The Tinetti balance scale was associated with H&Y stage, but the Tinetti gait scale was not. We assumed that this was because the H&Y stage is graded via laterality and postural imbalance and therefore cannot reflect gait status.

The TMT has been widely used to predict falls in the elderly.⁷ We found that cutoff values of 14 points on the Tinetti balance scale and 10 points on the gait scale showed the most accurate sensitivity and specificity for predicting falls among patients with PD. Previous reports also suggested that the optimal cutoff value of the Tinetti balance scale for predicting falling was 10 points among the elderly,^{16,17} but for patients with PD, the cutoff total score for the original TMT was 20 points, which is lower than our result.⁸ The Tinetti gait scale and balance scale showed acceptable accuracy for predicting falling in patients with PD in this study.

Our study has several limitations. First, the sam-

ple size was not large; therefore, the estimated power was low (55–62%). Further research will be needed to confirm the effect size of our study finding. Second, we did not enroll patients in an advanced stage of PD owing to their falling risk; therefore, interreliability for each item was low. Third, our study targeted only movement experts, not therapists or nurses.

In conclusion, our study demonstrated that the Korean version of the TMT has acceptable reliability and validity as an assessment tool for gait and balance among patients with PD. It is also useful for predicting falling among patients with PD using cut-off values of 14 on the balance scale and 10 on the gait scale.

Supplementary Materials

The online-only Data Supplement is available with this article at <https://doi.org/10.14802/jmd.17058>.

Conflicts of Interest

The authors have no financial conflicts of interest.

Acknowledgments

This research was funded by the Korean Movement Disorder Society (KMDS) as part of the “2016 Academic Research Funding” project.

REFERENCES

- Horak FB, Mancini M. Objective biomarkers of balance and gait for Parkinson's disease using body-worn sensors. *Mov Disord* 2013;28:1544-1551.
- Zhao YJ, Wee HL, Chan YH, Seah SH, Au WL, Lau PN, et al. Progression of Parkinson's disease as evaluated by Hoehn and Yahr stage transition times. *Mov Disord* 2010; 25:710-716.
- Movement Disorder Society Task Force on Rating Scales for Parkinson's Disease. The Unified Parkinson's Disease Rating Scale (UPDRS): status and recommendations. *Mov Disord* 2003;18:738-750.
- Perlmutter JS. Assessment of Parkinson disease manifestations. *Curr Protoc Neurosci* 2009;Chapter 10:Unit10.1.
- Qutubuddin AA, Pegg PO, Cifu DX, Brown R, McNamee S, Carne W. Validating the Berg Balance Scale for patients with Parkinson's disease: a key to rehabilitation evaluation. *Arch Phys Med Rehabil* 2005;86:789-792.
- Tinetti ME, Williams TF, Mayewski R. Fall risk index for elderly patients based on number of chronic disabilities. *Am J Med* 1986;80:429-434.
- Panzer VP, Wakefield DB, Hall CB, Wolfson LI. Mobility assessment: sensitivity and specificity of measurement sets in older adults. *Arch Phys Med Rehabil* 2011;92:905-912.
- Kegelmeyer DA, Kloos AD, Thomas KM, Kostyk SK. Reliability and validity of the Tinetti Mobility Test for individuals with Parkinson disease. *Phys Ther* 2007;87:1369-1378.
- An SH, Lee DG, Lee YB, Lee GC. Inter-rater? Absolute reliability and concurrent validity of Tinetti-gait scale (Korean version) in stroke patients. *J Korean Soc Phys Med* 2014;9: 201-211.
- Hughes AJ, Daniel SE, Blankson S, Lees AJ. A clinicopathologic study of 100 cases of Parkinson's disease. *Arch Neurol* 1993;50:140-148.
- Chiu AY, Au-Yeung SS, Lo SK. A comparison of four functional tests in discriminating fallers from non-fallers in older people. *Disabil Rehabil* 2003;25:45-50.
- McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)* 2012;22:276-282.
- Cipriani-Dacko LM, Innerst D, Johannsen J, Rude V. Interrater reliability of the Tinetti Balance Scores in novice and experienced physical therapy clinicians. *Arch Phys Med Rehabil* 1997;78:1160-1164.
- Yücel SD, Şahin F, Doğu B, Şahin T, Kuran B, Gürsakal S. Reliability and validity of the Turkish version of the Performance-Oriented Mobility Assessment I. *European Review of Aging and Physical Activity* 2012;9:149-159.
- Shore WS, deLateur BJ, Kuhlemeier KV, Imteyaz H, Rose G, Williams MA. A comparison of gait assessment methods: Tinetti and GAITRite electronic walkway. *J Am Geriatr Soc* 2005;53:2044-2045.
- Faber MJ, Bosscher RJ, van Wieringen PC. Clinimetric properties of the performance-oriented mobility assessment. *Phys Ther* 2006;86:944-954.
- Sterke CS, Huisman SL, van Beeck EF, Looman CW, van der Cammen TJ. Is the Tinetti Performance Oriented Mobility Assessment (POMA) a feasible and valid predictor of short-term fall risk in nursing home residents with dementia? *Int Psychogeriatr* 2010;22:254-263.