

Supplementary Table 6. KPPQ-Arabic Interrater agreement between first and second rater

	Rater A	Rater B	Kappa value	p value	ICC	95% CI
KPPQ item						
Q. 1: Pain around joints (musculoskeletal)	21 (45.7)	21 (45.7)	1	<0.001		
Q. 2: Pain related to internal organ	7 (15.2)	7 (15.2)	0.815	<0.001		
Q. 3: Lower abdominal pain	16 (34.8)	15 (32.6)	0.936	<0.001		
Q. 4: Pain deep within the body	7 (15.2)	6 (13.0)	0.903	<0.001		
Q. 5: Dyskinetic pain	7 (15.2)	8 (17.4)	0.912	<0.001		
Q. 6: "Off" dystonia in a region	16 (34.8)	16 (34.8)	1	<0.001		
Q. 7: Generalized "off" period pain	9 (19.6)	9 (19.6)	1	<0.001		
Q. 8: PLM or RLS-associated pain	15 (32.6)	14 (30.4)	0.935	<0.001		
Q. 9: Pain while turning in bed	13 (28.3)	14 (30.4)	0.934	<0.001		
Q. 10: Pain when chewing	5 (10.9)	4 (8.7)	0.807	<0.001		
Q. 11: Pain due to grinding teeth	6 (13.0)	6 (13.0)	1	<0.001		
Q. 12: Burning mouth syndrome	0 (0)	0 (0)	-	-		
Q. 13: Burning pain in the limbs	8 (17.4)	8 (17.4)	1	<0.001		
Q. 14: Shooting pain/pins and needles	12 (26.1)	12 (26.1)	1	<0.001		
KPPQ-Arabic total sore	4 (2–8)	4 (2–8)	-	-	0.996	0.991–0.998

Values are presented as *n* (%) or median (interquartile range).

KPPQ, King's Parkinson's Disease Pain Questionnaire; ICC, intraclass correlation coefficients; CI, confidence interval; PLM, periodic limb movements; RLS, restless legs syndrome.